Opioid Addiction Treatment Overview & Community Collaboration

Calvert County Behavioral Health Recovery Rapid Response & Mobile Crisis Team



Drew Fuller, MD, MPH
Medical Director




Anyone is Vulnerable



The Great Threat






72,000 People Died from Overdose in 2017






Calvert County – 10th Highest Death Rate in State




Leading Cause of Death Under Age 49


Real People – Parents & Children in S. Maryland













Heather Charnley 10/15/82 - 5/3/15



Overdoses 61



Deaths 14



24/7 MD Crisis Hotline: 1-800-422-0009 Hope4Calvert.org















1 Overdose Death in Calvert County Every

13 Days

Overdose Deaths in Calvert County 2019

Age
23
27
50
31
50
31
28
38
31
33
39
32
44
53
36
26
42
41

Average Age

34 yo

Young people with children are DYING DAILY



Grandparents as Primary Caregivers



_

Morphine = 1X Fentanyl =100X Carfentanil =10,000X

Lethal Dose





Lethal doses of heroin compared to "synthetic" opioids.

New Hampshire State Police Forensic Lab

'Death Pill': Fentanyl Disguised As Other Drugs Linked To Spike In US Overdoses











Addiction – Treatable Chronic Medical Disease



Biochemical and Structural Changes


Non-Opioid-Dependent and Opioid-Dependent Brain Images



PET scan images show changes in brain function caused by opioid dependence. The lack of red in the opioid-dependent brain shows a reduction in brain function in these regions.

Reprinted by permission of Nature Publishing Group: Neuropsychopharmacology. 1997;16:174-182.

Addiction: A Biopsychosocial Illness



Pediatric Development Disorder

Use often starts in teenage years

- Adverse Childhood Experiences
 - High ACEs Scores





In Deep





Lifestyle Disease?




80% Due to Lifestyle Choices

Diabetes Type 2



Relapse rates in chronic diseases




High Death Rate After Overdose & Narcan (1st responder)

9.9% Mortality (1 year)



Addiction and Suicide



Indiana Outbreak of HIV

The New york Times

Rural Indiana Struggles to Contend With H.I.V. Outbreak



Indiana HIV outbreak, hepatitis C epidemic sparks CDC alert

"This is one of the worst documented outbreaks of HIV among IV users in the past two decades."

Dr. Jonathan Mermin. director of the National Center for HIV/AIDS, Vir...

Indiana Reports More HIV Cases in Outbreak


Health officials confirm 120 HIV cases and 10 preliminary positive cases tied to Scott County

Associated Press

April 18, 2015

THE WALL STREET JOURNAL. HIV outbreak in Indiana grows


Hepatitis C



Age Distribution of All Hepatitis C Cases (Confirmed Acute and Confirmed/Probable Chronic) by Sex – New York State (Excluding NYC), 2014

Higher Death Rate <u>after</u> Hospital Admission (if no MAT)


Drug-related death rate per 1000




White S et al. Drugs-Related Death Soon after Hospital-Discharge among Drug Treatment Clients in Scotland: Record Linkage, Validation, and Investigation of Risk-Factors. PLoS One. 2015; 10(11): e0141073

Recently incarcerated people are over 40 times more likely to die from an opioid overdose

Number of opioid overdose deaths per 100,000 recently incarcerated people in North Carolina compared to rate among the general population in North Carolina









Which Treatments Work Best?


Treatment Effectiveness




Treatment Effectiveness



OAT Treatment Stability




Treatments Impact Different Brain Regions




Detox Does NOT Work

Opioid Detoxification Ineffective



Counseling Far More Effective with MAT



Evidence Based Treatment

"Access to medication – assisted treatment can mean [the] difference between life or death."


Michael Botticelli, October 23, 2014

Director, White House Office of National Drug Control Policy


Buprenorphine – Gold Standard, FDA Approved Treatment

- FDA Approval 38 years
 - 1981 for Pain
 - 2002 for MAT
- Safe- Very Low OD risk


Long Effect (24-36 hours)



Buprenorphine Binds 1000x Stronger Than Heroin




Partial Agonist — Highly Safe




Usually Once or Twice a day




Decreased Death Rate 80%



Impact Greater Than Most Medical Treatments



Comprehensive Approach



Broad Benefits of Treatment

- Get their lives back "Feel Normal Again"
- Better Parents
- Higher Employment
- Engaged in community
- Up to 80% decrease in mortality
- Decreased Crime 80%
- Decreased assoc illness
- 70-80% remain in recover
- Help Others

Myths & Misconceptions

"Trade one addiction for another"

Trade a life of chaos and harm for stable, safe & highly effective care

"they are just getting high"

Buprenorphine is not a desirable drug of abuse and can cause withdrawal

"They are going to sell the medication"

Diversion happens when treatment gaps exist. **Harm Reduction**. Attempt at self management

National Academy of Medicine



CONCLUSIONS OF THE NATIONAL ACADEMIES COMMITTEE

1. Opioid use disorder is a treatable chronic brain disease.

Opioid use disorder is a treatable chronic brain disease resulting from the changes in neural structure and function that are caused over time by repeated opioid use. The behavioral and social contexts are critically important both to its development and treatment. Stopping opioid misuse is extremely difficult. Medications are intended to normalize brain structure and function.

2. FDA-approved medications to treat opioid use disorder are effective and save lives.


FDA-approved medications to treat opioid use disorder—methadone, buprenorphine, and extended-release naltrexone—are effective and save lives. The most appropriate medication varies by individual and may change over time. To stem the opioid crisis, it is critical for all FDA-approved options to be available for all people with opioid use disorder. At the same time, as with all medical disorders, continued research on new medications, approaches, and formulations that will expand the options for patients is needed.

The National Academics of SCIENCES - ENGINEERING - MEDICHE CONSENSUS STUDY REPORT MEDICATIONS FOR OPIOID USE DISORDER SAVE LIVES

National Support for MAT



















Standard of Care


Medication-Assisted Treatment (MAT) is the standard of care for treating opioid use disorders:

- U.S. Dept. of Health & Human Services (1997)
 National Institute on Drug Abuse (2014, 2018)
 U.S. Surgeon General (2018)
 - Substance Abuse & Mental Health Services Administration (2005, 2018)
 - National Academy of Sciences, Engineering & Medicine (2019)
 - World Health Organization (2004)
 - Centers for Disease Control & Prevention (2002)
 - American Medical Association (2017)
 - American Psychiatric Association (2017)
 - American Society of Addiction Medicine (2015)
 - American Academy of Addiction Psychiatry
 - American College of Obstetricians & Gynecologists (2016)
 - National Association of Drug Court Professionals (2013, 2015)






Fewer than 20% of people with Opioid Use Disorder in Treatment




May be as low as 1/10 people with Opioid Use Disorder in Treatment



Window of Readiness Can Close Quickly



Put an end to stigma







Diversion & Harm Reduction

 Mostly used for self treatment or withdrawal avoidance (97%)*

Harm Reduction

- (Safer than any other opioid)
- Increase likely of later treatment
- Not a great drug of abuse
 - Slow onset
 - Blocks additional opioids
 - Precipitates withdrawal



Calvert County Help Lead State & Nation


Mobile Crisis Team (Opioid Response)

- October 7am-11pm
 - Goal 24/7/365 (TBA)

- PEER, Counselor, RN, Doctor/Nurse Practitioner (Live and Telehealth)
- Overdose and Crisis Response




Clinical Treatment Area



Mobile Crisis Team


Doctor/Nurse Practitioner Nurse



Licensed Counselor



PEER Recovery Specialist



"Meet Them Where They Are"



Hospital Collaboration

- ED and HD communication
- PEER Recovery Program
- MCT can be called to ER
- Quick Referral & Follow Up
 - Same day / Next Day
 - 7 Days a week



Community Partner Collaborations



EMS & Law Enforcement Collaboration

- Best Patient Engagement & Outcomes
- Decreased turn around times
- Decreased Repeat Calls
- Optimizes resource use



Activation of Mobile Crisis Team


911

CONTACT RECOVERY RAPID RESPONSE AT

877-467-5628

7 DAYS A WEEK

People Die Waiting for Treatment



Peles et al. J Addict Med 2013;7: 177-182

Behavioral Health Rapid Response

- Call Line 7AM-11PM
- Clinical Access 7 days a week
- Walk-in Access
- 4 Location in the County
 - Prince Frederick
 - Barstow (HQ)
 - Lusby
 - Chesapeake Beach

CONTACT RECOVERY RAPID RESPONSE AT

877-467-5628

7 DAYS A WEEK

Calvert County Health Dept at Chesapeake Beach





Calvert County Health Dept at Chesapeake Beach

- Open M-F 9-5
- Medically Assist Treatment
- Counseling
 - Individual
 - Group
- Intake Screening





"Treating addiction allows me to help individuals realize their hopes and dreams."

Christina Delos Reyes, MD, FASAM
Psychiatrist
ASAM member since 1995

Read My Story

Share Your Story

What If Everything Changed? We Could End OD



The Better Option

Opioid Addiction



MAT & Recovery



Progress Through Recovery


Team Effort




Their Destiny



Our Citizens Deserve the Most Effective Treatments







CALVERT COUNTY BEHAVIORAL HEALTH

RECOVERY RAPID RESPONSE

877-467-5628 7 DAYS A WEEK

Contacts

Drew Fuller, MD, MPH

Medical Director
Drew.Fuller@Maryland.gov
410-989-8181

Jason Phelan, BA

Supervisor of Crisis Response Coordination

jason.phelan@maryland.gov

Denise Dickerson, MS, LCADC, LCPC

Clinical Supervisor Rapid Response denise.dickerson@maryland.gov

CONTACT RECOVERY RAPID RESPONSE AT

877-467-5628

7 DAYS A WEEK